

144(6): New Spin Connection Resonance Structures

Define: $R_{\mu}^a = \frac{m}{e} r_{\mu}^a$ — (1)

Let: $\underline{A}_{orb}^a = \frac{\partial R^a}{\partial t} + c \underline{\nabla} R^a + c \omega_{ob}^a \underline{R}^b - c R^b \underline{\omega}^a$ — (2)

$\frac{1}{c} \underline{A}_{spi}^a = \underline{\nabla} \times \underline{R}^a - \underline{\omega}^a \times \underline{R}^b$ — (3)

From Cartan and Evans identities:

$\underline{\nabla} \cdot \underline{A}_{spi}^a = 0$ — (4)

$\underline{\nabla} \times \underline{A}_{orb}^a + \frac{1}{c} \frac{\partial \underline{A}_{spi}^a}{\partial t} = 0$ — (5)

$\underline{\nabla} \cdot \underline{A}_{orb}^a = \underline{J}^a$ — (6)

$\underline{\nabla} \times \underline{A}_{spi}^a - \frac{1}{c} \frac{\partial \underline{A}_{orb}^a}{\partial t} = \underline{J}^a$ — (7)

Eqs. (2) and (3), viewed as eqs. (4) to (7), produce various spin connection resonance

in: $R_{\mu}^a = (R^a, -R^a)$ — (8)
 $= \frac{m}{e} r_{\mu}^a$

This means that there is a special resonance property of \underline{A}_{spi}^a and \underline{A}_{orb}^a .